

# First Circum-Pacific Record of *Dimorphinites* (Ammonoidea, Upper Bajocian) in the Precordillera of Northern Chile

S.R. Fernandez-Lopez<sup>1\*</sup>, G.B. Chong Diaz<sup>2</sup>

1. Departamento de Paleontología, Facultad de Ciencias Geológicas and Instituto de Geología Económica, 28040-Madrid, Spain (E-mail: sixto@geo.ucm.es)

2. Departamento de Ciencias Geológicas, Universidad Católica del Norte, Antofagasta, Chile (E-mail: gchong@ucn.cl)

New ammonite fossil-assemblages containing morphoceratids have been discovered in the Quebrada San Pedro area, northern Chile, Region de Antofagasta, Comuna Sierra Gorda, approximately 40 km ESE of Sierra Gorda town and 75 km south of Calama city. The outcrop is located 1 km north of the composite section described as Quebrada de San Pedro and 10 km south of the classical localities of Caracoles Mining District previously studied by diverse authors (Gottsche, 1878; Steinmann, 1881; Möricke, 1894; Westermann, 1967, 1981; Hillebrandt, 1970, 1973, 1977, 2001; Westermann and Riccardi, 1979, 1980; Jensen and Quinzio, 1979, 1981; Gröschke and Hillebrandt, 1985; Hillebrandt et al., 1986; Gröschke et al., 1988; Riccardi et al., 1990a, b; Riccardi and Westermann, 1991a,b; Fernandez-Lopez et al., 1994; Gröschke and Hillebrandt, 1994).

In the Caracoles-Quebrada de San Pedro area, overlying unconformably Triassic volcanic and metamorphic rocks, there is a succession of Jurassic, fossiliferous deposits belonging to the Caracoles Group

(Harrington, 1961; Garcia, 1967; Montaña, 1976; Marinovic and Garcia, 1999; Ardill et al., 1998). The lowest marine deposits are brown, calcareous sandstones and fine-grained conglomerates belonging to the Coronado Formation. This lithostratigraphic unit is followed by yellow and brown shales with intercalations of limestones and calcareous sandstones, of the Torcazas Formation, developed in shallow marine settings. In overlying levels, there are yellow and green shales with intercalations of limestones, belonging to the Mina Chica Formation, developed in offshore settings. The lithostratigraphic position of the new ammonite fossil-assemblages is the top of the Torcazas Formation, below the Mina Chica Formation. In this locality, the last beds of the Torcazas Formation correspond to a centimetric interval of grey wackestone to packstone limestones containing resedimented specimens of Upper Bajocian ammonites, without traces of reelaboration. These ammonites have been collected by one of the authors (F.L.) during several field studies, between 2006 and 2010.






Age	Lithostratigraphy	Column	Lithology	Setting
Middle Jurassic	Caracoles Formation	 18 m	Grey and blue limestones with intercalations of green shales	Shallow marine
	Mina Chica Formation	 75 m	Yellow and green shales with intercalations of limestones	Offshore
	Torcazas Formation	 100 m	Yellow and brown shales with intercalations of limestones and calcareous sandstones	Shallow marine
	Coronado Formation	 40 m	Brown calcareous sandstones and fine-grained conglomerates	Shallow marine
Trias.	Agua Dulce Formation		Violet and red silicified volcanic rocks, including lava flows, tuffs and breccias	Volcanic

Fig. 1 Lithostratigraphy and lithology of study area

Among the ammonoids of these fossil-assemblages, Sphaeroceratidae are dominant, being *Megasphaeroceras* [M+m] the most common ammonite genus. Stephanoceratidae, Perisphinctidae and Oppeliidae are scarce, but *Cadomites* [M]-*Polyploectites* [m], *Vermisphinctes* [m]-*Prorsisphinctes* [M] and *Oxyerites* [M]-*Paroecotraustes* [m] are well represented. Morphoceratidae, Lissoceratidae and Strigoceratidae are very scarce, respectively represented by *Dimor-*

*phinites* [M]-*Vigoricerias* [m], *Lissoceras* [M]-*Micro-lissoceras* [m] and *Strigoceras* [M]. Besides Ammonitina, Phylloceratina and Lytoceratina are extremely scarce, but they occur. Among the morphoceratids of this stratigraphical interval, macroconchs of *Dimorphinites dimorphus* (D'Orbigny) and microconchs of *Vigoricerias defranciai* (D'Orbigny) have been identified. Since *Dimorphinites dimorphus* (D'Orbigny) [M+m] has only been known from the

Parkinsoni Zone, Upper Bajocian, of Europe, North Africa and south part of the Transcaucasus (cf. Dietze and Bernt, 2009), this standard zone is now extended to the SE Pacific. Some associated taxa, such as *Vermisphinctes* [m]-*Prorsisphinctes* [M], *Strigoceras* [M] and *Oxycerites* [M]-*Paroecotraustes* [m] also indicate the uppermost Bajocian age and Tethyan affinities.

**Key words:** Jurassic; Marine stratigraphy; Palaeobiogeography; South America

## References:

- Ardill J., Flint S., Chong G., et al. Sequence stratigraphy of the Mesozoic Domeyko Basin, northern Chile. *Journal of the Geological Society, London*, 1998, 155: 71-88.
- Dietze V., Bernt E. Erstnachweis der Gattung *Dimorphinites* Buckman, 1923 (Morphoceratidae) aus dem Ober-Bajocium (Mittlerer Jura) Süddeutschlands. *Jahreshefte der Gesellschaft für Naturkunde in Württemberg*, 2009, 165: 5-11.
- Fernandez-Lopez S., Chong Diaz G., Quinzio Sinn L.A., et al. The Upper Bajocian and Bathonian in the Cordillera de Domeyko, North-Chilean Pre-cordillera: Sedimentological and biostratigraphical results. *Geobios, M.S.*, 1994, 17: 187-201.
- Garcia F. Geología del Norte Grande de Chile. Sociedad Geológica de Chile. Simposio sobre el Geosinclinal Andino, 1967: 1-138.
- Gottsche C. Über jurassische Versteinerungen aus der argentinischen Cordillere. *Palaeontographica*, 1878, supplement III, Lieferung II, Heft 2: 1-50.
- Gröschke M., von Hillebrandt A. Trias und Jura in der mittleren Cordillera Domeyko von Chile (221 30' -241 30'). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 1985, 170: 129-166.
- Gröschke M., von Hillebrandt A. The Bathonian in Northern Chile. *Geobios, M.S.*, 1994, 17:255-264.
- Gröschke M., von Hillebrandt A., Prinz P., et al. Marine Mesozoic Paleogeography in Northern Chile between 211-261 S. *Lecture Notes Earth Sciences*, 1988, 17: 105-117.
- Harrington H.L. Geology of parts of Antofagasta and Atacama Provinces, Northern Chile. *The American Association of Petroleum Geologists Bulletin*, 1961, 45: 169-197.
- von Hillebrandt A. Zur Biostratigraphie und Ammoniten-Fauna des südamerikanischen Jura (insbes. Chile). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 1970, 136: 166-211.
- von Hillebrandt A. Neue Ergebnisse über den Jura in Chile und Argentinien. *Münstersche Forschungen zur Geologie und Palaontologie*, 1973 (31/32): 167-199.
- von Hillebrandt A. Ammoniten aus dem Bajocien (Jura) von Chile (Südamerika). *Neue Arten der Gattung* *Stephanoceras* und *Domeykoceras* n. gen. (Stephanoceratidae). *Mitteilungen der Bayerischen Staatssammlung für Palaontologie und Historische Geologie*, 1977, 17: 35-69.
- von Hillebrandt A. Ammonite stratigraphy of the Bajocian in Northern Chile. *Hantkeniana*, 2001, 3: 49-87.
- von Hillebrandt A., Gröschke M., Prinz P., et al. Marines Mesozoikum in Nordchile zwischen 211 und 261S. *Berliner Geowissenschaftliche Abhandlungen*, 1986, 66: 169-190.
- Jensen A., Quinzio L.A. Geología del Área de Pampa Elvira y contribución al conocimiento del Jurásico marino entre los 231 00' y 231 30' Latitud Sur y los 681 45' y 691 03' Longitud Oeste. II Región de Antofagasta, Chile. *Memoria Universidad Católica del Norte, Dpto. Geociencias, Antofagasta*, 1979: 1-141.
- Jensen A., Quinzio L.A. Hallazgo de Batoniano y descripción de facies sedimentarias marinas del Dogger en Quebrada San Pedro, Caracoles, II Región-Antofagasta (Chile). *Estudios geológicos*, 1981, 37: 209-214.
- Marinovic N., García M. Hoja Pampa Unión, región de Antofagasta. 1:100,000. Servicio Nacional de Geología y Minería. Mapas Geológicos, 1999, N Mapa: M33
- Möricke W. Versteinerungen des Lias und Unteroolith von Chile. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie*, 1894, 9: 1-100.
- Montaño J.M. Estudio geológico de la zona de Caracoles y áreas vecinas, con énfasis en el Sistema Jurásico, provincia de Antofagasta, II Región, Chile. Tesis, Santiago: Universidad de Chile, Departamento de Geología, 1976: 182.
- Riccardi A.C., Westermann G.E.G. Middle Jurassic ammonite fauna and biochronology of the Argentine-Chilean Andes. Part III: Bathonian-Callovian Eurycephalitinae, Stephanocerataceae. *Palaeontographica*, 1991a, 216: 1-110.
- Riccardi A.C., Westermann G.E.G. Middle Jurassic ammonite fauna and biochronology of the Argentine-Chilean Andes. Part IV: Bathonian-Callovian Reineckeidae. *Palaeontographica*, 1991b, 216: 111-145.
- Riccardi A.C., Westermann G.E.G., Elmi S. The Middle Jurassic Bathonian-Callovian Ammonite Zones of the Argentine-Chilean Andes. *Geobios*, 1990a, 22: 553-597.
- Riccardi A.C., Westermann G.E.G., Damborenea S.E. Middle Jurassic of South America and Antarctic Peninsula. *Newsletters on Stratigraphy*, 1990b, 21: 105-128.
- Steinmann G. Zur Kenntnis der Jura und Kreideformation von Caracoles (Bolivia). *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie*, 1881, 49: 239-301.
- Westermann G.E.G. Sucesión de ammonites del Jurá-

sico medio en Antofagasta, Atacama, Mendoza y Neuquén. Revista de la Asociación Geológica Argentina, 1967, 22: 65-73.

Westermann G.E.G. Ammonite biochronology and biogeography of the Circum Pacific Middle Jurassic. The Systematics Association, 1981, 18: 459-498.

Westermann G.E.G., Riccardi A.C. Middle Jurassic

ammonoid fauna and biochronology of the Argentine-Chilean Andes, Part. II, Bajocian Stephanocerataceae. Palaeontographica, 1979, 164: 85-188.

Westermann G.E.G., Riccardi A.C. The Upper Bajocian ammonite *Strenoceras* in Chile: First circum-Pacific record of the Subfurcatum Zone: Newsletters on Stratigraphy, 1980, 9: 19-29.




地质学报

SHORT PAPERS FOR THE 8<sup>TH</sup> INTERNATIONAL CONGRESS ON THE JURASSIC SYSTEM

2010年 第17卷 特刊

中国地质大学(北京)  
北京 大学

# 地质学报



SHORT PAPERS FOR THE 8<sup>TH</sup> INTERNATIONAL  
CONGRESS ON THE JURASSIC SYSTEM

Marine and non-marine Jurassic

**Special Issue**  
2010 Vol.17